

This article was downloaded by:[Rosier, L.]
On: 21 November 2007
Access Details: [subscription number 783432256]
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



International Journal of Control

Publication details, including instructions for authors and subscription information:
<http://www.informaworld.com/smpp/title~content=t713393989>

Output feedback stabilization of a clamped-free beam

P. Le Gall^a, C. Prieur^b; L. Rosier^a

^a Institut Élie Cartan, UMR 7502 UHP/CNRS/INRIA, B.P. 239, F-54506

Vandoeuvre-lès-Nancy Cedex, France

^b LAAS-CNRS, 31077 Toulouse, France

First Published on: 01 June 2007

To cite this Article: Gall, P. Le, Prieur, C. and Rosier, L. (2007) 'Output feedback stabilization of a clamped-free beam', *International Journal of Control*, 80:8, 1201 - 1216

To link to this article: DOI: 10.1080/00207170601165097

URL: <http://dx.doi.org/10.1080/00207170601165097>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Output feedback stabilization of a clamped-free beam

P. LE GALL[†], C. PRIEUR[‡] and L. ROSIER^{*†}

[†]Institut Élie Cartan, UMR 7502 UHP/CNRS/INRIA, B.P. 239, F-54506
Vandœuvre-lès-Nancy Cedex, France

[‡]LAAS-CNRS, 7 avenue du colonel Roche,
31077 Toulouse, France

(Received 17 July 2006; in final form 5 September 2006)

We consider a Euler–Bernoulli beam, clamped at one extremity and free at the other, to which are attached a piezoelectric actuator and a collocated sensor touching the clamped extremity. We provide an output feedback law and characterize the sensor/actuator lengths for which the strong stabilization holds. Finally, we prove that the energy decreases to zero in a polynomial way for almost all lengths, and in an exponential way for lengths admitting a certain coprime factorization.

1. Introduction

The Euler–Bernoulli equation provides a very commonly used model for the dynamic behaviour of a flexible beam, when the cross-sectional dimensions of the beam are small in comparison to its length. In the last decade, a lot of papers were devoted to the pointwise feedback stabilization of a beam; see e.g. Chen *et al.* (1987), Conrad (1990), Lagnese and Leugering (1991a, b), Morgül (1992), Rebarber (1995), Liu and Zheng (1999), Ammari and Tucsnak (2000) and de Queiroz *et al.* (2000) or to the modelling of piezoceramic actuation of beams (Crawley and Anderson 1990, Banks and Smith 1992) or to the controllability of such systems (Crépeau and Prieur (to appear), Tucsnak 1996). The piezoceramic actuation of beams or plates is investigated numerically and experimentally in Destuynder *et al.* (1992), Banks *et al.* (1993) and Destuynder (1999).

In this paper, we consider the control problem modelling the vibrations of a Euler–Bernoulli beam which is subject to the action of an attached piezoelectric actuator. If we assume that the beam is clamped at one end and free at the other, and that the

actuator is touching the clamped end, then we obtain the system

$$\begin{aligned}w_{tt}(x, t) + w_{xxxx}(x, t) &= -h(t) \frac{d\delta_\xi}{dx}(x), \\w(0, t) = w_x(0, t) = w_{xx}(\pi, t) = w_{xxx}(\pi, t) &= 0, \\w(x, 0) = w^0(x), w_t(x, 0) &= w^1(x).\end{aligned}$$

In the above equations, $x \in (0, \pi)$ is the spatial coordinate, t is time, w stands for the transverse deflection of the beam, $w_t = \partial w / \partial t$, etc., 0 and ξ are the coordinates of the extremities of the actuator ($0 < \xi \leq \pi$), δ_y is the Dirac measure at the point y , and h stands for the control input (see figure 1).

We are interested in the stabilization of the above system, the control h being expressed as a function of the output $w_x(0, t) - w_x(\xi, t) = -w_x(\xi, t)$. This corresponds to the situation where the output comes from a piezoelectric sensor located on the same interval $(0, \xi)$ as the actuator. Formal computations on the variations of the energy

$$E(w, w_t) = \int_0^\pi (|w_{xx}|^2 + |w_t|^2) dx$$

lead to take

$$h(t) := -Kw_{xt}(\xi, t)$$

*Corresponding author. Email: rosier@iecn.u-nancy.fr




Figure 1. Clamped-free beam with collocated sensor/actuator.

where K is some number whose range will be specified later. In short, we are interested in the stability properties of the closed-loop system

$$w_{tt}(x, t) + w_{xxxx}(x, t) = Kw_{xt}(\xi, t) \frac{d\delta_\xi}{dx}(x), \quad (1)$$

$$w(0, t) = w_x(0, t) = w_{xx}(\pi, t) = w_{xxx}(\pi, t) = 0, \quad (2)$$

$$w(x, 0) = w^0(x), w_t(x, 0) = w^1(x). \quad (3)$$

The main goal of the paper is to characterize the lengths of the sensor/actuator leading to “good” stability properties for (1)–(3), i.e., with an exponential (or polynomial) energy decay rate.

The study of the controllability properties of a PDE is often considered as the first step towards the control theory of that PDE. The exact controllability of a beam hinged at both ends and subject to the action of a piezoelectric actuator is characterized in function of the location of the actuator in Tucsnak (1996). It is shown in that paper that the space of controllable states is related to Diophantine approximation properties for the extremities of the actuator. A similar study is performed in Crépeau and Prieur (to appear) in the more difficult framework of a clamped-free beam. The stabilization of a beam hinged at both ends and subject to a pointwise force has been deeply studied in Ammari and Tucsnak (2000). The method of proof consists of deducing the decay estimates from observability inequalities for the associated free problem via sharp trace regularity results.

When we compare our work with Ammari and Tucsnak (2000) we notice that: (i) the boundary conditions here, where only an asymptotic expansion of the eigenvalues for the free evolution of (1)–(3) is known, are more difficult to handle than the ones in Ammari and Tucsnak (2000) (where e.g. the eigenvectors for the free evolution reduce to sin functions); (ii) the method of proof used in Ammari and Tucsnak (2000) cannot be applied

here due to an additional difficulty. The key observed quantity, namely the integral term $\int_0^T |w_{xt}(\xi, t)|^2 dt$, is of the same order as $\|(w^0, w^1)\|_{H^3 \times H^1}^2$ for the free evolution ($K=0$), whereas it is of the same order as $\|(w^0, w^1)\|_{H^2 \times H^0}^2$ for the controlled system ($K > 0$). Thus, a spectral analysis resting on the free evolution as in Ammari and Tucsnak (2000) cannot be performed here.

To obtain our decay estimates, we follow the frequency-domain approach described in Liu and Zheng (1999) and based on a sharp estimation of a resolvent on the imaginary axis. More precisely, we combine the multiplier method to a careful computation of the value at $x = \pi$ of the solution to (1)–(3). Thanks to that calculation, we obtain an explicit polynomial decay estimate valid for almost every value of ξ , and an exponential decay estimate when ξ admits a certain coprime factorization.

The paper is outlined as follows. The well-posedness of (1)–(3) in the energy space is investigated in §2. The lengths ξ of the actuator for which the strong stability holds are characterized in §3. The last section contains the main results of the paper, namely Theorem 3 (resp., Theorem 4) which provides an exponential (resp., polynomial) energy decay rate.

2. Well-posedness of (1)–(3)

The well-posedness of (1)–(3) in the energy space is obtained in this section as a direct application of the classical semi-group theory. To specify the operator, we have to see for which function w the r.h.s. of (1) belongs to $L^2(0, \pi)$. Let us introduce some notations. If w is any function in $H^1(0, \xi) \cap H^1(\xi, \pi)$, we define $\{w_x\} \in L^2(0, \pi)$ by

$$\{w_x\}(x) := \begin{cases} w_x^{D(0, \xi)}(x) & \text{if } x \in (0, \xi), \\ w_x^{D(\xi, \pi)}(x) & \text{if } x \in (\xi, \pi), \end{cases}$$

where $w_x^{D'(0, \xi)}$ (resp. $w_x^{D'(\xi, \pi)}$) denotes the distributional derivative $\partial w / \partial x$ in $D'(0, \xi)$ (resp. in $D'(\xi, \pi)$). We also set $[w]_\xi := w(\xi^+) - w(\xi^-)$. It follows that

$$w_x = \{w_x\} + [w]_\xi \delta_\xi \quad \text{in } D'(0, \pi).$$

Assume now that $w, v \in H^2(0, \pi)$, and define $u \in D'(0, \pi)$ by $u := -w_{xxxx} + K v_x(\xi) d\delta_\xi / dx$. If $u \in L^2(0, \pi)$, then the restriction of w_{xxxx} to each of the intervals $(0, \xi)$ and (ξ, π) has to be a square-integrable function; hence $w \in H^4(0, \xi) \cap H^4(\xi, \pi)$. Calculating w_{xxx} , w_{xxxx} and u , we obtain

$$\begin{aligned} w_{xxx} &= \{w_{xxx}\} + [w_{xx}]_\xi \delta_\xi, \\ w_{xxxx} &= \{w_{xxxx}\} + [w_{xxx}]_\xi \delta_\xi + [w_{xx}]_\xi \frac{d}{dx} \delta_\xi, \end{aligned}$$

and

$$\begin{aligned} u &= -\{w_{xxxx}\} - [w_{xxx}]_\xi \delta_\xi - [w_{xx}]_\xi \frac{d}{dx} \delta_\xi \\ &\quad + K v_x(\xi) \frac{d}{dx} \delta_\xi. \end{aligned} \tag{4}$$

Then $u \in L^2(0, \pi)$ provided that all the coefficients in front of the Dirac measures vanish, i.e.,

$$K v_x(\xi) = [w_{xx}]_\xi$$

and

$$[w_{xxxx}]_\xi = 0.$$

We are now in a position to define the operator associated with (1)–(3).

Let $V = \{w \in H^2(0, \pi) \mid w(0) = w'(\pi) = 0\}$ (where $' = d/dx$) $(w_1, w_2)_V = \int_0^\pi w_1' \overline{w_2'} dx$, and $H = L^2(0, \pi)$, $(v_1, v_2)_H = \int_0^\pi v_1 \overline{v_2} dx$. Then $\mathcal{H} = V \times H$, endowed with the usual product norm, is a (complex) Hilbert space. If we introduce $v := w_t$ and define the operator \mathcal{A} with domain

$$\begin{aligned} \mathcal{D}(\mathcal{A}) &= \{z = (w, v) \mid (w, v) \in H^2(0, \pi)^2, \\ &\quad w \in H^4(0, \xi) \cap H^4(\xi, \pi), \\ &\quad w(0) = w_x(0) = w_{xx}(\pi) = w_{xxx}(\pi) = 0, \quad v(0) = v_x(0) = 0, \\ &\quad K v_x(\xi) = [w_{xx}]_\xi, [w_{xxxx}]_\xi = 0\} \end{aligned}$$

by

$$\mathcal{A}z = \left(v, -w_{xxxx} + K v_x(\xi) \frac{d}{dx} \delta_\xi \right) = (v, -\{w_{xxxx}\}), \tag{5}$$

then we see that the closed-loop system (1)–(3) may be interpreted as the initial value problem for the abstract first-order evolution equation in \mathcal{H}

$$\begin{cases} \frac{dz}{dt} = \mathcal{A}z, & t > 0 \\ z(0) = (w^0, w^1). \end{cases}$$

The first result in this paper is the following one.

Theorem 1: *If $K \geq 0$, then \mathcal{A} generates a C_0 -semigroup $(e^{t\mathcal{A}})_{t \geq 0}$ of contractions in \mathcal{H} .*

Proof: Obviously, $\mathcal{D}(\mathcal{A})$ is dense in \mathcal{H} . According to a classical result (see e.g. Liu and Zheng (1999, [Theorem 1.2.4]), the theorem is proved if we show that \mathcal{A} is dissipative and that $0 \in \rho(\mathcal{A})$, the resolvent set of \mathcal{A} . Let us begin with the dissipativity property.

Lemma 1: *For any $z = (w, v) \in \mathcal{D}(\mathcal{A})$ we have that*

$$(\mathcal{A}z, z)_\mathcal{H} = 2i \operatorname{Im} \left(\int_0^\pi v_{xx} \overline{w_{xx}} dx \right) - K |v_x(\xi)|^2. \tag{6}$$

In particular, $\operatorname{Re}(\mathcal{A}z, z)_\mathcal{H} = -K |v_x(\xi)|^2 \leq 0$, i.e., \mathcal{A} is dissipative.

Proof of Lemma 1: Pick any pair of functions $(w, v) \in \mathcal{D}(\mathcal{A})$. Then

$$(\mathcal{A}(w, v), (w, v))_\mathcal{H} = \int_0^\pi v_{xx} \overline{w_{xx}} dx - \int_0^\pi \{w_{xxxx}\} \overline{v} dx.$$

After some integrations by parts on the intervals $(0, \xi)$ and (ξ, π) , we obtain that

$$\begin{aligned} - \int_0^\pi \{w_{xxxx}\} \overline{v} dx &= - \int_0^\xi \{w_{xxxx}\} \overline{v} dx - \int_\xi^\pi \{w_{xxxx}\} \overline{v} dx \\ &= - \int_0^\pi w_{xx} \overline{v_{xx}} dx + [w_{xx} \overline{v_x}]_{x=0}^\xi + [w_{xx} \overline{v_x}]_{x=\xi}^\pi \\ &= - \int_0^\pi w_{xx} \overline{v_{xx}} dx - [w_{xx}]_\xi \overline{v_x(\xi)}. \end{aligned}$$

Hence

$$\begin{aligned} (\mathcal{A}(w, v), (w, v))_\mathcal{H} &= \int_0^\pi (v_{xx} \overline{w_{xx}} - w_{xx} \overline{v_{xx}}) dx - [w_{xx}]_\xi \overline{v_x(\xi)} \\ &= 2i \operatorname{Im} \left(\int_0^\pi v_{xx} \overline{w_{xx}} dx \right) - K |v_x(\xi)|^2. \end{aligned}$$

We now proceed to the study of \mathcal{A}^{-1} . The following result holds true. □

Proposition 1: $0 \in \rho(\mathcal{A})$.

Proof: We have to prove that the operator $\mathcal{A}: \mathcal{D}(\mathcal{A}) \rightarrow \mathcal{H}$ is one-to-one and onto, and that its inverse $\mathcal{A}^{-1}: \mathcal{H} \rightarrow \mathcal{H}$ is continuous. Let a pair $(f, g) \in \mathcal{H}$ be given, and let us investigate the equation $\mathcal{A}(w, v) = (f, g)$, where (w, v) has to be found in $\mathcal{D}(\mathcal{A})$. We have to solve the system

$$\begin{cases} v = f \\ -\{w_{xxxx}\} = g \end{cases}$$

supplemented by adequate boundary conditions. To eliminate g , we introduce the (unique) solution $\tilde{w} \in H^4(0, \pi)$ of the following elliptic problem

$$\begin{cases} -\tilde{w}_{xxxx} = g \\ \tilde{w}(0) = \tilde{w}_x(0) = \tilde{w}_{xx}(\pi) = \tilde{w}_{xxx}(\pi) = 0. \end{cases}$$

Setting $w = \tilde{w} + \hat{w}$ and $C := K\nu_x(\xi) = Kf_x(\xi)$, we have to solve

$$\{\hat{w}_{xxxx}\} = 0 \tag{7}$$

$$\hat{w}(0) = \hat{w}_x(0) = \hat{w}_{xx}(\pi) = \hat{w}_{xxx}(\pi) = 0 \tag{8}$$

$$[\hat{w}_{xx}]_\xi = C \tag{9}$$

$$[\hat{w}_{xxx}]_\xi = 0 \tag{10}$$

where \hat{w} is to be found in $H^2(0, \pi) \cap H^4(0, \xi) \cap H^4(\xi, \pi)$. In particular,

$$[\hat{w}]_\xi = [\hat{w}_x]_\xi = 0. \tag{11}$$

We infer from (7) and (8) that there exist some constants a_2, a_3, b_0 , and b_1 such that

$$\hat{w}(x) = \begin{cases} a_2x^2 + a_3x^3 & \text{if } 0 < x < \xi, \\ b_0 + b_1(x - \pi) & \text{if } \xi < x < \pi. \end{cases}$$

It follows from (10) that $a_3=0$. Equation (9) gives $-2a_2 = C$; hence $a_2 = -C/2$. Finally, it is easily seen that b_0 and b_1 are uniquely determined by the following system of linear equations (coming from (11))

$$\begin{cases} b_0 + b_1(\xi - \pi) = a_2\xi^2, \\ b_1 = 2a_2\xi. \end{cases}$$

This proves the existence and uniqueness of \hat{w} , and the existence and uniqueness of a pair $(w, v) \in \mathcal{D}(\mathcal{A})$ such that $\mathcal{A}(w, v) = (f, g)$. To see that $0 \in \rho(\mathcal{A})$, it remains to prove that the map $\mathcal{A}^{-1}: \mathcal{H} \rightarrow \mathcal{H}$ is continuous. Let $(w, v) = \mathcal{A}^{-1}(f, g)$. Then $\|v\|_H = \|f\|_H$, $\|\tilde{w}\|_{H^4(0, \pi)} \leq \text{Const}\|g\|_H$, $\|\hat{w}\|_V \leq \text{Const}|C| \leq \text{Const}\|f\|_V$. Therefore,

$$\|(w, v)\|_{\mathcal{H}} \leq \text{Const}\|(f, g)\|_{\mathcal{H}}.$$

This completes the proofs of Proposition 1 and of Theorem 1. \square

Proposition 2: *The operator $\mathcal{A}^{-1}: \mathcal{H} \rightarrow \mathcal{H}$ is compact.*

Proof: Since

$$\|\mathcal{A}^{-1}(f, g)\|_{\mathcal{D}(\mathcal{A})} = \|\mathcal{A}^{-1}(f, g)\|_{\mathcal{H}} + \|(f, g)\|_{\mathcal{H}},$$

we see that \mathcal{A}^{-1} is continuous from \mathcal{D} into $\mathcal{D}(\mathcal{A})$, therefore it is sufficient to prove that the embedding $\mathcal{D}(\mathcal{A}) \rightarrow \mathcal{H}$ is compact. Let $((w^n, v^n))_{n \geq 0}$ be any bounded sequence in $\mathcal{D}(\mathcal{A})$. We have to prove that a subsequence $((w^{n'}, v^{n'}))$ converges (strongly) in \mathcal{H} . As $\|v^n\|_{H^2(0, \pi)} \leq \text{Const}$., there exist a subsequence $(v^{n'})$ and a function $v \in H = L^2(0, \pi)$ such that $v^{n'} \rightarrow v$ in H . On the other hand, $\|w^n\|_{H^2(0, \pi)} + \|\{w^n_{xxxx}\}\|_{L^2(0, \pi)} \leq \text{Const}$., so

$$\|w^{n'}\|_{H^4(0, \xi)} + \|w^{n'}\|_{H^4(\xi, \pi)} \leq \text{Const}.$$

Extracting if needed a subsequence again denoted by $((w^{n'}, v^{n'}))$, we infer that there exists a function $w \in V$ such that $w^{n'} \rightarrow w$ in $H^2(0, \pi)$, and $w^{n'} \rightarrow w$ in $H^4(0, \xi)$ and in $H^4(\xi, \pi)$. It follows that $w^{n'} \rightarrow w$ in $H^2(0, \xi)$ and in $H^2(\xi, \pi)$. We conclude that $w^{n'} \rightarrow w$ in V . \square

3. Strong stability

3.1 Free evolution

In this section we recall some useful facts about the free evolution of (1)–(3); i.e., when $K=0$. Thus we consider the homogeneous Cauchy problem

$$\phi_{tt} + \phi_{xxxx} = 0, \tag{12}$$

$$\phi(0, t) = \phi_x(0, t) = \phi_{xx}(\pi, t) = \phi_{xxx}(\pi, t) = 0, \tag{13}$$

$$\phi(., 0) = \phi^0, \quad \phi_t(., 0) = \phi^1. \tag{14}$$

Let $A: \mathcal{D}(A) \rightarrow L^2(0, \pi)$ be the operator with domain

$$\mathcal{D}(A) := \{\phi \in H^4(0, \pi) \mid \phi(0) = \phi_x(0) = \phi_{xx}(\pi) = \phi_{xxx}(\pi) = 0\}$$

and defined by $A\phi = \phi_{xxxx}$. Obviously, A^{-1} is a compact symmetric operator on $L^2(0, \pi)$, hence there exists a countable orthonormal basis of $L^2(0, \pi)$ constituted of eigenvectors of A^{-1} (and of A). The following result [(Crepeau and Prieur (to appear), Lemma 2.1] provides useful results about the eigenvectors of A .

Proposition 3: The $L^2(0, \pi)$ -normalized eigenfunctions of A are the functions $(\psi_k)_{k \geq 1}$ defined by

$$\psi_k(x) = \frac{1}{\sqrt{\pi}} \left\{ \cos(\alpha_k x) - \cosh(\alpha_k x) + \mu_k (\sinh(\alpha_k x) - \sin(\alpha_k x)) \right\}, \quad (15)$$

where α_k is the k th positive root of

$$1 + \cos(\alpha_k \pi) \cosh(\alpha_k \pi) = 0, \quad (16)$$

and

$$\mu_k = \frac{\cos(\alpha_k \pi) + \cosh(\alpha_k \pi)}{\sin(\alpha_k \pi) + \sinh(\alpha_k \pi)},$$

and the eigenvalue associated with ψ_k is $\lambda_k = |\alpha_k|^4$. Moreover, we have as $k \rightarrow +\infty$

$$\alpha_k = k - \frac{1}{2} + (-1)^{k+1} \frac{2e^{\pi/2}}{\pi} e^{-\pi k} + o(e^{-\pi k}), \quad (17)$$

$$\mu_k = 1 + 2(-1)^k e^{-\alpha_k \pi} + o(e^{-\alpha_k \pi}), \quad (18)$$

and

$$\begin{aligned} & -\sinh(\alpha_k \rho) + \mu_k \cosh(\alpha_k \rho) \\ &= \begin{cases} e^{-\alpha_k \rho} + o(e^{-\alpha_k \rho}) & \text{if } 0 < \rho < \pi/2, \\ (-1)^k e^{-\alpha_k(\pi-\rho)} + o(e^{-\alpha_k(\pi-\rho)}) & \text{if } \pi/2 < \rho < \pi. \end{cases} \end{aligned} \quad (19)$$

It is easily seen that if $\phi^0 = \sum_{k \geq 1} \phi_k^0 \psi_k$, and $\phi^1 = \sum_{k \geq 1} \phi_k^1 \psi_k$, then the solution $\phi = \phi(x, t)$ of (12)–(14) reads

$$\phi(x, t) = \sum_{k=1}^{+\infty} \left(\phi_k^0 \cos(|\alpha_k|^2 t) + \frac{\phi_k^1}{|\alpha_k|^2} \sin(|\alpha_k|^2 t) \right) \psi_k(x). \quad (20)$$

3.2 Statement of the strong stability result

For any $k \geq 1$ let

$$\mathcal{S}_k := \{ \xi \in (0, \pi] \mid \psi'_k(\xi) = 0 \}, \quad (21)$$

and set

$$\mathcal{S} := \cup_{k \geq 1} \mathcal{S}_k.$$

Definition 1: We say that (1)–(3) is strongly stable in \mathcal{H} , if for any $(w^0, w^1) \in \mathcal{H}$ we have that

$$E(w(t), w_t(t)) = \|(w(t), w_t(t))\|_{\mathcal{H}}^2 \rightarrow 0$$

as $t \rightarrow +\infty$.

The following theorem provides a characterization of the lengths of the sensor/actuator for which the strong stability holds.

Theorem 2: The system (1)–(3) is strongly stable in \mathcal{H} if and only if $K > 0$ and $\xi \notin \mathcal{S}$.

Proof: First, it follows from Lemma 1 that for any $(w^0, w^1) \in \mathcal{D}(A)$ and for any $t \geq 0$,

$$E(w(T), v(T)) - E(w^0, w^1) = -2K \int_0^T |v_x(\xi, t)|^2 dt. \quad (22)$$

Thus, the condition $K > 0$ is needed for the energy to decrease. On the other hand, if $\xi \in \mathcal{S}_k$ for some $k \geq 1$, then any state of the form $(\phi^0, \phi^1) = (\phi_k^0 \psi_k, \phi_k^1 \psi_k)$ gives rise to a solution of (1)–(3) whose energy does not tend to 0. Thus $K > 0$ together with $\xi \notin \mathcal{S}$ constitutes a necessary condition for (1)–(3) to be strongly stable. Let us now check that this condition is also sufficient. We need the following result, which extends Proposition 1. \square

Proposition 4 $i\mathbb{R} \subset \rho(\mathcal{A})$.

Proof of Proposition 4: Since \mathcal{A}^{-1} is compact, the spectrum of \mathcal{A} is constituted only of eigenvalues, so to prove the claim it is sufficient to check that for any $\beta \in \mathbb{R}^*$ the equation

$$(i\beta - \mathcal{A})(w, v) = (0, 0), \quad (w, v) \in \mathcal{D}(\mathcal{A}) \quad (23)$$

admits only the trivial solution $(w, v) = (0, 0)$. Using (23) and (6), we obtain that

$$\begin{aligned} 0 &= ((i\beta - \mathcal{A})(w, v), (w, v))_{\mathcal{H}} \\ &= i \left(\beta \|(w, v)\|_{\mathcal{H}}^2 - 2 \operatorname{Im} \int_0^\pi v_{xx} \overline{w_{xx}} dx \right) + K |v_x(\xi)|^2, \end{aligned}$$

hence

$$[w_{xx}]_{\xi} = K v_x(\xi) = 0$$

and $w \in H^4(0, \pi)$. On the other hand, (23) yields

$$\begin{cases} -\beta^2 w + w^{(4)} = 0 \\ w(0) = w'(0) = w''(\pi) = w'''(\pi) = 0, \end{cases}$$

so if $w \neq 0$, then $\beta^2 = \lambda_k$ and $w = C \psi_k$ for some $k \geq 1$, $C \in \mathbb{C}$. It follows that $0 = K v'(\xi) = i \beta K C \psi'_k(\xi)$; hence $C = 0$ (since $K \neq 0$ and $\xi \notin \mathcal{S}$), contradicting the assumption $w \neq 0$. Therefore $w = v = 0$.

Applying a classical stability theorem due to Arendt–Batty (1988), we infer from Proposition 4 that

the system (1)–(3) is strongly stable in \mathcal{H} . This completes the proof of Theorem 2. \square

Remark 1:

- (i) Proceeding as in Le Gall *et al.* (to appear), one may write a more elementary proof of Theorem 2 by using LaSalle’s invariance principle (see Ammari and Tucsnak (2000) for strong stability results obtained this way).
- (ii) If $\xi \in \mathcal{S}_k$ and $\xi \notin \mathcal{S}_{k'}$ for $k' \neq k$, then a strong stability result also holds true in the subspace $\mathcal{H}' := \text{Span}\{(\psi_k, 0), (0, \psi_k)\}^\perp$ of codimension 2.

3.3 Properties of the set \mathcal{S} of critical lengths

As the set \mathcal{S} plays a crucial role in all the stability results, we collect some of its properties in the following proposition.

Proposition 5: *The set \mathcal{S} is countable and dense in $(0, \pi]$.*

Proof: To prove that \mathcal{S} is (at most) countable, it is sufficient to prove that each set \mathcal{S}_k is finite. But $\mathcal{S}_k = (\psi'_k)^{-1}(0) \cap (0, \pi]$, and the function ψ'_k , which is analytic, has only a finite number of zeros in the interval $(0, \pi]$ if it is not identically null. To check the last property, we need first to establish Claim 1.

Claim 1 ($\mu_k \neq 1 \quad \forall k \geq 1$). Argue by contradiction. If the claim is false, then there exists some $k \geq 1$ with

$$\mu_k = \frac{\cos(\alpha_k \pi) + \cosh(\alpha_k \pi)}{\sin(\alpha_k \pi) + \sinh(\alpha_k \pi)} = 1. \tag{24}$$

Let $x = \cos(\alpha_k \pi)$. Using (16) and (24), we arrive to the equation

$$x - x^{-1} = \pm \sqrt{1 - x^2} + \sqrt{x^{-2} - 1}$$

whose solutions are easily found to be ± 1 . Now, (16) has no solution if $\cos(\alpha_k \pi) = \pm 1$. The claim is proved.

Derivating in (15) we obtain

$$\begin{aligned} \psi'_k(x) &= \pi^{-1/2} \left(-\alpha_k \sin(\alpha_k x) - \alpha_k \sinh(\alpha_k x) \right. \\ &\quad \left. + \mu_k (\alpha_k \cosh(\alpha_k x) - \alpha_k \cos(\alpha_k x)) \right) \\ &\sim \pi^{-1/2} \alpha_k (\mu_k \cosh(\alpha_k x) - \sinh(\alpha_k x)) \\ &\sim \frac{\pi^{-1/2} \alpha_k (\mu_k - 1) e^{\alpha_k x}}{2} \end{aligned}$$

as $x \rightarrow +\infty$, hence $\psi'_k \neq 0$. The proof that \mathcal{S} is at most countable is achieved.

Let us now check that \mathcal{S} is dense in $(0, \pi]$ (hence not finite). Applying Proposition 3, we obtain that

$$\psi'_k(x) = -\sqrt{2} \pi^{-1/2} \alpha_k \left(\sin\left(\alpha_k x + \frac{\pi}{4}\right) + O(e^{-\delta \alpha_k}) \right)$$

on each of the intervals $[0, \pi/2 - \varepsilon]$ and $[\pi/2 + \varepsilon, \pi]$ ($\varepsilon > 0$ being arbitrarily small), $\delta = \delta(\varepsilon) \in (0, \pi/2)$ denoting some appropriate constant. Applying the intermediate values theorem, we conclude that the function ψ'_k vanishes between any pair of successive extrema of the function $\sin(\alpha_k x + \pi/4)$ on each of the above intervals. The density of \mathcal{S} follows at once. \square

4. Energy decay rates

In this section we provide explicit energy decay rates in function of the length ξ of the sensor/actuator.

4.1 Exponential decay rate

An exponential decay rate is first derived when ξ/π is rational with

$$\frac{\xi}{\pi} \neq \frac{4k' + 3}{4k + 2}, \quad \forall k, k' \in \mathbb{Z}. \tag{25}$$

Theorem 3: *Let $\xi \notin \mathcal{S}$ be such that $\xi/\pi \in \mathbb{Q}$ and (25) holds true. Then*

$$\sup_{|\beta| \geq 1} \|(i\beta - \mathcal{A})^{-1}\| < \infty. \tag{26}$$

It follows that the semigroup $(e^{t\mathcal{A}})_{t \geq 0}$ is exponentially stable; i.e., there exist two constants $C > 0$ and $\delta > 0$ such that

$$\|e^{t\mathcal{A}} z_0\|_{\mathcal{H}} \leq C e^{-\delta t} \|z_0\|_{\mathcal{H}}$$

for any $z_0 = (w_0, v_0) \in \mathcal{H}$ and any $t \geq 0$.

Remark 2: It is easy to see that $\xi := \pi$ does not belong to \mathcal{S} , and that (25) is satisfied for this choice of ξ . Unfortunately, checking that $\xi \notin \mathcal{S}$ for a ξ fulfilling (25) seems to be a very hard task.

Proof of Theorem 3: The exponential stability of the semigroup $(e^{t\mathcal{A}})_{t \geq 0}$ is in fact equivalent to the resolvent estimate (26) by virtue of a well-known result due to Huang and Prüss, see Prüss (1984) and Huang (1985). Therefore, we shall focus on the proof of (26). As the usefulness of the condition (25) will appear quite far in the proof, we assume at the beginning of the proof

that ξ is any number in $(0, \pi]$. Let us argue by contradiction. If (26) is false, then there exist $\beta_n \in \mathbb{R}$, $(w_n, v_n) \in \mathcal{D}(\mathcal{A})$ for $n = 1, 2, \dots$ such that

$$\|(w_n, v_n)\|_{\mathcal{H}} = 1, \quad |\beta_n| \rightarrow +\infty \quad (27)$$

and if we set $(f_n, g_n) := (i\beta_n - \mathcal{A})(w_n, v_n)$,

$$(f_n, g_n) \rightarrow (0, 0) \quad \text{in } \mathcal{H}, \quad (28)$$

i.e.

$$i\beta_n w_n - v_n = f_n \rightarrow 0 \quad \text{in } V, \quad (29)$$

$$i\beta_n v_n + \{w_n^{(4)}\} = g_n \rightarrow 0 \quad \text{in } H. \quad (30)$$

Replacing if necessary β_n , w_n and v_n by $-\beta_n$, \bar{w}_n and \bar{v}_n , respectively, we may assume that $\beta_n > 0$ for all n . Our goal is to obtain a contradiction to (27). This will be done in four steps. In the first step, using the multiplier method we establish some estimates which show that we are done if $\beta_n w_n(\pi) \rightarrow 0$ as $n \rightarrow +\infty$. In the second step, we show that the couple $(w_n(\pi), w_n'(\pi))$ may be obtained as the solution of a 2×2 linear system by performing a direct integration of (29) and (30). In the two remaining steps we estimate the determinants involved in the resolution of that linear system. In what follows, the letters C, C', C'', \dots will denote positive constants which may vary from line to line.

Step 1 (Basic estimates by the multiplier method): We infer from (6) that

$$\begin{aligned} K|v_n'(\xi)|^2 &= \operatorname{Re}((i\beta_n - \mathcal{A})(w_n, v_n), (w_n, v_n))_{\mathcal{H}} \\ &= \operatorname{Re}((f_n, g_n), (w_n, v_n))_{\mathcal{H}}, \end{aligned} \quad (31)$$

hence, using (27) and (28),

$$|v_n'(\xi)|^2 \rightarrow 0 \quad \text{as } n \rightarrow +\infty. \quad (32)$$

On the other hand, (29) and (30) give as $n \rightarrow \infty$

$$i\beta_n \|w_n\|_V^2 - (v_n, w_n)_V \rightarrow 0, \quad (33)$$

$$i\beta_n \|v_n\|_H^2 + (\{w_n^{(4)}\}, v_n)_H \rightarrow 0. \quad (34)$$

Taking the difference of (33) and (34) we obtain

$$i\beta_n (\|w_n\|_V^2 - \|v_n\|_H^2) - \int_0^\pi v_n'' \bar{w}_n' dx - \int_0^\pi \{w_n^{(4)}\} \bar{v}_n dx \rightarrow 0$$

but

$$\operatorname{Im} \left(\int_0^\pi v_n'' \bar{w}_n' dx + \int_0^\pi \{w_n^{(4)}\} \bar{v}_n dx \right) = 0$$

by (6), hence

$$\beta_n (\|w_n\|_V^2 - \|v_n\|_H^2) \rightarrow 0. \quad (35)$$

Therefore, using (27) and (29) we conclude that

$$\lim_{n \rightarrow +\infty} \|w_n\|_V^2 = \lim_{n \rightarrow +\infty} \|v_n\|_H^2 = \lim_{n \rightarrow +\infty} \|\beta_n w_n\|_H^2 = \frac{1}{2}. \quad (36)$$

Eliminating v_n in (30) by using (29), we obtain

$$-\beta_n^2 w_n + \{w_n^{(4)}\} = g_n + i\beta_n f_n, \quad (37)$$

hence

$$(-\beta_n^2 w_n + \{w_n^{(4)}\}, qw_n')_H = (g_n + i\beta_n f_n, qw_n')_H \quad (38)$$

for any real function $q \in C^3([0, \pi])$.

It follows from (36) and a classical interpolation inequality (see e.g. Lions and Magenes (1972)) that for any $s \in [0, 2)$

$$w_n \rightarrow 0 \quad \text{in } H^s(0, \pi). \quad (39)$$

In particular,

$$(g_n, qw_n')_H \rightarrow 0. \quad (40)$$

On the other hand,

$$\int_0^\pi f_n q \bar{w}_n' dx = - \int_0^\pi (f_n q)' \bar{w}_n dx + f_n(\pi) q(\pi) \overline{w_n(\pi)} \quad (41)$$

but, using (29) and (36), we obtain

$$\beta_n \left| \int_0^\pi (f_n q)' \bar{w}_n dx \right| \leq C \|f_n\|_{H^1(0, \pi)} \|\beta_n w_n\|_H \rightarrow 0, \quad (42)$$

and

$$\beta_n f_n(\pi) q(\pi) \overline{w_n(\pi)} = o(\beta_n w_n(\pi)). \quad (43)$$

It follows that

$$(i\beta_n f_n, qw_n')_H = o(\beta_n w_n(\pi)) + o(1). \quad (44)$$

Let us now turn to the left hand side of (38). We have

$$2\text{Re}(-\beta_n^2 w_n, qw'_n)_H = \beta_n^2 \int_0^\pi q' |w_n|^2 dx - \beta_n^2 |w_n(\pi)|^2 q(\pi). \tag{45}$$

Integrating by parts in

$$\left(\{w_n^{(4)}\}, qw'_n \right)_H = \int_0^\xi w_n^{(4)} q \overline{w'_n} dx + \int_\xi^\pi w_n^{(4)} q \overline{w'_n} dx,$$

we arrive to

$$\begin{aligned} 2\text{Re}(\{w_n^{(4)}\}, qw'_n)_H &= 3 \int_0^\pi |w_n''|^2 q' dx + |w_n''(0)|^2 q(0) \\ &\quad - \int_0^\pi |w_n'|^2 q''' dx \\ &\quad + q(\xi) [|w_n''|^2]_\xi + 2\text{Re} \left(q'(\xi) \overline{w_n''(\xi)} [w_n'']_\xi \right) \\ &\quad + |w_n'(\pi)|^2 q''(\pi) \\ &= 3 \int_0^\pi |w_n''|^2 q' dx \\ &\quad + 2Kq(\xi) \text{Re}(v_n'(\xi) \overline{w_n''(\xi^+)}) + o(1) \end{aligned} \tag{46}$$

by (39) and (32), provided that $q(0) = 0$. Gathering together (40), (44), (45) and (46) with $q(x) = x$, we obtain

$$\begin{aligned} &3 \int_0^\pi |w_n''|^2 dx + \beta_n^2 \int_0^\pi |w_n|^2 dx \\ &= (\pi \beta_n \overline{w_n(\pi)} + o(1)) \beta_n w_n(\pi) \\ &\quad - 2K\xi \text{Re}(v_n'(\xi) \overline{w_n''(\xi^+)}) + o(1). \end{aligned} \tag{47}$$

The above equality may be simplified thanks to the next result.

Proposition 6:

$$|w_n''(\xi^+)|^2 = O(|\beta_n w_n(\pi)|^2) + o(1). \tag{48}$$

Proof: Multiplying each term in (37) by $q \overline{w'_n}$ where $q \in C^3([\xi, \pi])$, and integrating over (ξ, π) , we arrive to

$$-\beta_n^2 \int_\xi^\pi qw_n \overline{w'_n} dx + \int_\xi^\pi qw_n^{(4)} \overline{w'_n} dx = \int_\xi^\pi (g_n + i\beta_n f_n) q \overline{w'_n} dx.$$

Clearly

$$\left| \int_\xi^\pi g_n q \overline{w'_n} dx \right| \leq C \|g_n\|_H \|w'_n\|_H \rightarrow 0$$

and

$$\int_\xi^\pi i\beta_n f_n q \overline{w'_n} dx = -i \int_\xi^\pi (f_n q)' \beta_n \overline{w_n} dx + [if_n q \beta_n \overline{w_n}]_\xi^\pi,$$

so using again (29) and (36),

$$\left| \int_\xi^\pi (g_n + i\beta_n f_n) q \overline{w'_n} dx \right| \leq \frac{\xi}{4} (|\beta_n w_n(\xi)|^2 + |\beta_n w_n(\pi)|^2) + o(1).$$

On the other hand, we obtain after some integrations by parts that

$$2\text{Re} \int_\xi^\pi (-\beta_n^2) qw_n \overline{w'_n} dx = \beta_n^2 \left\{ \int_\xi^\pi q' |w_n|^2 dx - [q |w_n|^2]_\xi^\pi \right\}$$

and that

$$\begin{aligned} 2\text{Re} \int_\xi^\pi w_n^{(4)} q \overline{w'_n} dx &= 3 \int_\xi^\pi q' |w_n''|^2 dx - [|w_n''|^2 q]_\xi^\pi \\ &\quad - \int_\xi^\pi q''' |w_n'|^2 dx + [q'' |w_n'|^2]_\xi^\pi \\ &\quad - 2\text{Re} [w_n'' q' \overline{w'_n}]_\xi^\pi + 2\text{Re} [w_n''' q \overline{w'_n}]_\xi^\pi. \end{aligned}$$

Taking $q(x) = x$ and using the fact that $w_n''(\pi) = w_n'''(\pi) = 0$, we arrive to

$$\begin{aligned} &3 \int_\xi^\pi |w_n''|^2 dx + \beta_n^2 \int_\xi^\pi |w_n|^2 dx + \xi (|w_n''(\xi^+)|^2 \\ &\quad + \frac{1}{2} |\beta_n w_n(\xi)|^2) + 2\text{Re}((w_n''(\xi^+) - \xi w_n'''(\xi)) \overline{w_n'(\xi)}) \\ &= O(|\beta_n w_n(\pi)|^2) + o(1). \end{aligned} \tag{49}$$

The following claims are needed. □

Claim 2: $\beta_n w_n'(\xi) \rightarrow 0$ as $n \rightarrow \infty$.

Indeed, derivating once in (29) and evaluating at $x = \xi$, we infer that $i\beta_n w_n'(\xi) = v_n'(\xi) + f_n'(\xi) \rightarrow 0$, by (28) and (32).

Claim 3: $\|w_n\|_{H^4(\xi, \pi)} \leq C\beta_n$.

Indeed, it follows from (30) and (36) that

$$\|\beta_n^{-1} w_n^{(4)}\|_{L^2(\xi, \pi)} \leq \beta_n^{-1} \|g_n\|_{L^2(\xi, \pi)} + \|v_n\|_{L^2(\xi, \pi)} \leq C.$$

Since $\|w_n\|_V$ is bounded, the claim follows.

We infer from Claim 3 that $|w_n''(\xi^+)| + |w_n'''(\xi)| \leq C\beta_n$ which, combined to Claim 2, yields

$$\text{Re}((w_n''(\xi^+) - \xi w_n'''(\xi)) \overline{w_n'(\xi)}) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore, (49) yields

$$3 \int_{\xi}^{\pi} |w_n''|^2 dx + \beta_n^2 \int_{\xi}^{\pi} |w_n|^2 dx + \xi(|w_n''(\xi^+)|^2 + \frac{1}{2}|\beta_n w_n(\xi)|^2) = O(|\beta_n w_n(\xi)|^2) + o(1),$$

hence (48) follows. The proof of Proposition 6 is complete. \square

Using (32), (47) and (48), we arrive to

$$3 \int_0^{\pi} |w_n''|^2 dx + \beta_n^2 \int_0^{\pi} |w_n|^2 dx = O(|\beta_n w_n(\pi)|^2) + o(1). \tag{50}$$

Therefore, we obtain a contradiction with (36) if $\beta_n w_n(\pi) \rightarrow 0$. In the next steps, we show that $\beta_n w_n(\pi) \rightarrow 0$.

Step 2 Computation of $w_n(\pi)$: To compute $w_n(\pi)$ we integrate (37) on $(0, \pi)$. More precisely, setting $F_n := g_n + i\beta_n f_n$, we solve the system

$$\begin{cases} -\beta_n^2 w_n + \{w_n^{(4)}\} = F_n & \text{on } (0, \pi), \\ w_n(0) = w_n'(0) = w_n''(\pi) = w_n'''(\pi) = 0, \\ [w_n]_{\xi} = [w_n']_{\xi} = [w_n'']_{\xi} = 0, \\ [w_n'']_{\xi} = K v_n'(\xi) = K(i\beta_n w_n'(\xi) - f_n'(\xi)). \end{cases}$$

To simplify the notations we drop the subscript n in what follows. Let λ and μ be given numbers. We first solve the following (backwards) Cauchy Problem on (ξ, π)

$$(S_1) \quad \begin{cases} w^{(4)} - \beta^2 w = F & \text{in } (\xi, \pi), \\ w(\pi) = \lambda, w'(\pi) = \mu, w''(\pi) = w'''(\pi) = 0, \end{cases}$$

and next the following (backwards) Cauchy Problem on $(0, \xi)$

$$(S_2) \quad \begin{cases} w^{(4)} - \beta^2 w = F & \text{in } (0, \xi), \\ w(\xi^-) = w(\xi^+), w'(\xi^-) = w'(\xi^+), w''(\xi^-) = w''(\xi^+), \\ w''(\xi^-) = w''(\xi^+) - K(i\beta w'(\xi^+) - f'(\xi)). \end{cases}$$

Then λ and μ have to be chosen so that $w(0) = w'(0) = 0$.

The ODE $w^{(4)} - \beta^2 w = F$ may be written as

$$\begin{pmatrix} w \\ w' \\ w'' \\ w''' \end{pmatrix}' = M \begin{pmatrix} w \\ w' \\ w'' \\ w''' \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ 0 \\ F \end{pmatrix}, \quad \text{with}$$

$$M = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ \beta^2 & 0 & 0 & 0 \end{pmatrix}.$$

Easy computations show that

$$e^{xM} = \begin{pmatrix} a & \beta^{-2} a''' & \beta^{-2} a'' & \beta^{-2} a' \\ a' & a & \beta^{-2} a''' & \beta^{-2} a'' \\ a'' & a' & a & \beta^{-2} a''' \\ a''' & a'' & a' & a \end{pmatrix}, \tag{51}$$

where $a(x) := (\cos(bx) + \cosh(bx))/2$ and $b = \sqrt{\beta}$. An integration of (S_1) yields

$$\begin{pmatrix} w \\ w' \\ w'' \\ w''' \end{pmatrix}(x) = e^{(x-\pi)M} \begin{pmatrix} \lambda \\ \mu \\ 0 \\ 0 \end{pmatrix} + \int_{\pi}^x e^{(x-y)M} \begin{pmatrix} 0 \\ 0 \\ 0 \\ F(y) \end{pmatrix} dy, \quad x \in (\xi, \pi),$$

hence

$$\begin{pmatrix} w \\ w' \\ w'' \\ w''' \end{pmatrix}(\xi^+) = e^{(\xi-\pi)M} \begin{pmatrix} \lambda \\ \mu \\ 0 \\ 0 \end{pmatrix} + \int_{\pi}^{\xi} e^{(\xi-y)M} \begin{pmatrix} 0 \\ 0 \\ 0 \\ F(y) \end{pmatrix} dy. \tag{52}$$

On the other hand, an integration of (S_2) leads to

$$\begin{pmatrix} w \\ w' \\ w'' \\ w''' \end{pmatrix}(x) = e^{(x-\xi)M} \begin{pmatrix} w \\ w' \\ w'' \\ w''' \end{pmatrix}(\xi^-) + \int_{\xi}^x e^{(x-y)M} \begin{pmatrix} 0 \\ 0 \\ 0 \\ F(y) \end{pmatrix} dy. \tag{53}$$

Gathering (52) and (53), we obtain

$$\begin{pmatrix} w \\ w' \\ w'' \\ w''' \end{pmatrix}(0) = e^{-\xi M} \begin{pmatrix} w \\ w' \\ w'' \\ w''' \end{pmatrix}(\xi^+) + \begin{pmatrix} 0 \\ 0 \\ -[w'']_{\xi} \\ 0 \end{pmatrix} + \int_{\xi}^0 e^{-yM} \begin{pmatrix} 0 \\ 0 \\ 0 \\ F(y) \end{pmatrix} dy = e^{-\pi M} \begin{pmatrix} \lambda \\ \mu \\ 0 \\ 0 \end{pmatrix} + \int_{\pi}^0 e^{-yM} \begin{pmatrix} 0 \\ 0 \\ 0 \\ F(y) \end{pmatrix} dy + e^{-\xi M} \begin{pmatrix} 0 \\ 0 \\ -K(i\beta w'(\xi^+) - f(\xi)) \\ 0 \end{pmatrix}. \tag{54}$$

By (51) and (52),

$$w'(\xi^+) = a'(\xi - \pi)\lambda + a(\xi - \pi)\mu + \int_{\pi}^{\xi} \beta^{-2} a''(\xi - y)F(y) dy,$$

hence, keeping only the two first equations in (54), we arrive to

$$\begin{pmatrix} w(0) \\ w'(0) \end{pmatrix} = \begin{pmatrix} a(-\pi) & \beta^{-2} a'''(-\pi) \\ a'(-\pi) & a(-\pi) \end{pmatrix} \begin{pmatrix} \lambda \\ \mu \end{pmatrix} + \int_{\pi}^0 \begin{pmatrix} \beta^{-2} a'(-y) \\ \beta^{-2} a''(-y) \end{pmatrix} F(y) dy + \begin{pmatrix} \beta^{-2} a''(-\xi) \\ \beta^{-2} a'''(-\xi) \end{pmatrix} \times (Kf'(\xi) - iK\beta(a'(\xi - \pi)\lambda + a(\xi - \pi)\mu + \int_{\pi}^{\xi} \beta^{-2} a''(\xi - y)F(y) dy)).$$

Therefore, $w(0) = w'(0) = 0$ if and only if

$$\begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{pmatrix} \begin{pmatrix} \lambda \\ \mu \end{pmatrix} = \begin{pmatrix} c_1 \\ c_2 \end{pmatrix},$$

where

$$\begin{aligned} m_{11} &= a(-\pi) - iK\beta^{-1} a''(-\xi) a'(\xi - \pi) \\ m_{12} &= \beta^{-2} a'''(-\pi) - iK\beta^{-1} a''(-\xi) a(\xi - \pi) \\ m_{21} &= a'(-\pi) - iK\beta^{-1} a'''(-\xi) a'(\xi - \pi) \\ m_{22} &= a(-\pi) - iK\beta^{-1} a''(-\xi) a(\xi - \pi) \\ c_1 &= - \int_{\pi}^0 \beta^{-2} a'(-y)F(y) dy + K\beta^{-2} a''(-\xi) \\ &\quad \times \left(-f'(\xi) + i\beta \int_{\pi}^{\xi} \beta^{-2} a''(\xi - y)F(y) dy \right) \\ c_2 &= - \int_{\pi}^0 \beta^{-2} a''(-y)F(y) dy + K\beta^{-2} a'''(-\xi) \\ &\quad \times \left(-f'(\xi) + i\beta \int_{\pi}^{\xi} \beta^{-2} a''(\xi - y)F(y) dy \right). \end{aligned}$$

Letting

$$\begin{aligned} N &= c_1 m_{22} - c_2 m_{12} \\ D &= m_{11} m_{22} - m_{12} m_{21} \end{aligned}$$

we obtain by Cramer rule $\lambda = N/D$. In the next step we show that $|D| \geq Ce^{b\pi}$ if the number ξ/π is rational and fulfills (25).

Step 3 Estimation of D : Substituting the expressions of m_{11}, m_{12}, m_{21} and m_{22} into D we obtain

$$\begin{aligned} D &= (a(-\pi) - iK\beta^{-1} a''(-\xi) a'(\xi - \pi))(a(-\pi) - iK\beta^{-1} a'''(-\xi) a(\xi - \pi)) - (\beta^{-2} a'''(-\pi) - iK\beta^{-1} a''(-\xi) a(\xi - \pi))(a'(-\pi) - iK\beta^{-1} a'''(-\xi) a'(\xi - \pi)) \\ &= a(-\pi)^2 - \beta^{-2} a'''(-\pi) a'(-\pi) - iK\beta^{-1} \{ a''(-\xi) a'(\xi - \pi) a(-\pi) + a(-\pi) a'''(-\xi) a(\xi - \pi) - a'(-\xi) a(\xi - \pi) a'(-\pi) - \beta^{-2} a'''(-\pi) a'''(-\xi) a'(\xi - \pi) \} \\ &=: D_1 + iD_2. \end{aligned}$$

Substituting the values of a and of its derivatives in D_1 we obtain

$$\begin{aligned} D_1 &= \left(\frac{1}{2} (\cos b\pi + \cosh b\pi) \right)^2 - \beta^{-2} \frac{b^3}{2} (-\sin b\pi - \sinh b\pi) \\ &\quad \times \frac{b}{2} (\sin b\pi - \sinh b\pi) \\ &= \frac{1}{4} (\cos^2 b\pi + \cosh^2 b\pi + 2 \cos b\pi \cosh b\pi + \sin^2 b\pi - \sinh^2 b\pi) \\ &= \frac{1}{2} (1 + \cos b\pi \cosh b\pi). \end{aligned} \tag{55}$$

Let us now turn to the estimation of D_2 .

$$D_2 = -K\beta^{-1} \{ a'(\xi - \pi)(a''(-\xi)a(-\pi) - \beta^{-2}a'''(-\pi)a'''(-\xi)) + a(\xi - \pi)(a(-\pi)a'''(-\xi) - a''(-\xi)a'(-\pi)) \}.$$

To estimate D_2 we introduce the functions $G, H: [0, \pi] \rightarrow \mathbb{R}$ defined by

$$G(y) = a''(-\xi)a(-y) - \beta^{-2}a'''(-y)a'''(-\xi) \\ H(y) = a(-y)a'''(-\xi) - a''(-\xi)a'(-y).$$

Then the following result holds.

Lemma 2:

$$G(y) = \frac{b^2}{4} \left(\cosh b(y - \xi) + \frac{e^{b\xi}}{2} (\cos by - \sin by) - \frac{e^{by}}{2} (\cos b\xi + \sin b\xi) + O(1) \right), \quad (56)$$

$$H(y) = \frac{b^3}{4} \left(\sinh b(y - \xi) - \frac{e^{b\xi}}{2} (\cos by + \sin by) - \frac{e^{by}}{2} (\cos b\xi + \sin b\xi) + O(b) \right), \quad (57)$$

where $O(1)$ denotes a term which is bounded in y, ξ , and b .

Proof of Lemma 2: Expanding a in the expression of G , we obtain

$$G(y) = \frac{b^2}{4} \{ (-\cos b\xi + \cosh b\xi)(\cos by + \cosh by) - (\sin by + \sinh by)(\sin b\xi + \sinh b\xi) \} \\ = \frac{b^2}{4} \{ \cosh by \cosh b\xi - \sinh by \sinh b\xi + \frac{1}{2}(e^{b\xi} + e^{-b\xi}) \cos by - \frac{1}{2}(e^{by} + e^{-by}) \cos b\xi - \frac{1}{2}(e^{by} - e^{-by}) \sin b\xi - \frac{1}{2}(e^{b\xi} - e^{-b\xi}) \sin by - \cos b\xi \cos by - \sin by \sin b\xi \} \\ = \frac{b^2}{4} \left\{ \cosh b(y - \xi) + \frac{e^{b\xi}}{2} (\cos by - \sin by) - \frac{e^{by}}{2} (\cos b\xi + \sin b\xi) + O(1) \right\}.$$

Thus (56) is proved. (57) may be obtained by noticing that $H(y) = G'(y)$. \square

Applying Lemma 2 with $y = \pi$, we obtain

$$D_2 = -K\beta^{-1} (a'(\xi - \pi)G(\pi) + a(\xi - \pi)H(\pi)) \\ = -K\beta^{-1} \left\{ \frac{b}{2} \left(-\sin b(\xi - \pi) + \frac{e^{b(\xi - \pi)} - e^{b(\pi - \xi)}}{2} \right) \times \frac{b^2}{4} \left(\frac{e^{b(\pi - \xi)}}{2} + \frac{e^{b\xi}}{2} (\cos b\pi - \sin b\pi) - \frac{e^{b\pi}}{2} (\cos b\xi + \sin b\xi) + O(1) \right) + \frac{1}{2} \left(\cos b(\xi - \pi) + \frac{e^{b(\xi - \pi)} + e^{b(\pi - \xi)}}{2} \right) \times \frac{b^3}{4} \left(\frac{e^{b(\pi - \xi)}}{2} - \frac{e^{b\xi}}{2} (\cos b\pi + \sin b\pi) - \frac{e^{b\pi}}{2} (\cos b\xi + \sin b\xi) + O(1) \right) \right\}.$$

Expanding D_2 , we notice that the coefficients in front of the exponential terms $e^{b(2\pi - \xi)}$ and $e^{2b(\pi - \xi)}$ vanish, so that the leading exponential term is $e^{b\pi}$, and the remaining exponential terms read $e^{b(\pi - \xi)}$, $e^{b(2\xi - \pi)}$, $e^{b\xi}$, e^0 , and $e^{b(\xi - \pi)}$. We conclude that if $\xi < \pi$, then

$$D_2 = \frac{Kb}{16} (\cos b\pi + (\cos b\xi + \sin b\xi) \times (\cos b(\xi - \pi) - \sin b(\xi - \pi))e^{b\pi} + O(e^{b(\pi - \delta)})) \quad (58)$$

for some $\delta > 0$, and if $\xi = \pi$, then

$$D_2 = \frac{Kb}{4} ((\cos b\pi + \sin b\pi)e^{b\pi} + O(b)). \quad (59)$$

It is clear that $|D_1 + iD_2| \geq Ce^{b\pi}$ for b large enough if $\xi = \pi$, for $(\cos b\pi)^2 + (\cos b\pi + \sin b\pi)^2 \geq C > 0$ for all $b \in \mathbb{R}$.

Let us assume from now on that $\xi < \pi$, and that the number ξ/π is rational and admits a coprime factorization $\xi/\pi = p/q$, with $p, q \in \mathbb{N}^*$, $0 < p < q$, and let us introduce the functions

$$c(b) := \cos b\pi + (\cos b\xi + \sin b\xi) \times (\cos b(\xi - \pi) - \sin b(\xi - \pi)), \quad (60)$$

$$h(b) := (\cos b\pi)^2 + (c(b))^2. \quad (61)$$

Lemma 3: $\inf_{b \in \mathbb{R}} |h(b)| > 0$ if and only if p/q is not of the form $(4k' + 3)/(4k + 2)$, $k, k' \in \mathbb{Z}$.

Proof of Lemma 3: Since h is $2q$ -periodic, we only have to characterize the rational numbers p/q for which $h(b) > 0$ for all $b \in \mathbb{R}$. As h is the sum of two squared terms, we may restrict ourselves to the b 's

for which $\cos b\pi = 0$, namely $b = (1/2) + k$, $k \in \mathbb{Z}$. Then $h(b) = 4 \cos^2((1/2 + k)\xi - \pi/4) \cos^2((1/2 + k) \times (\xi - \pi) + \pi/4)$, and a straightforward calculation shows that $h(b) = 0$ if and only if p/q is of the form $(4k' + 3)/(4k + 2)$, $k, k' \in \mathbb{Z}$. \square

Applying Lemma 3, we infer that as $b \rightarrow +\infty$

$$\begin{aligned} |D|^2 &= D_1^2 + D_2^2 \\ &\geq Ch(b)e^{2b\pi} + O(e^{b(2\pi-\delta)}) \\ &\geq C'e^{2b\pi} \end{aligned}$$

provided that the condition (25) is fulfilled.

Step 4 Estimation of N : We have to bound the quantity $N = c_1 m_{22} - c_2 m_{12}$. Letting

$$Z_1 := - \int_{\pi}^0 \beta^{-2} a'(-y) F(y) dy$$

$$Z_2 := - \int_{\pi}^0 \beta^{-2} a''(-y) F(y) dy$$

and
$$Z_3 := i\beta^{-1} \int_{\pi}^{\xi} a''(\xi - y) F(y) dy - f'(\xi),$$

we obtain that

$$c_1 = Z_1 + K\beta^{-2} a''(-\xi) Z_3, \quad c_2 = Z_2 + K\beta^{-2} a'''(-\xi) Z_3,$$

hence

$$\begin{aligned} N &= (Z_1 + K\beta^{-2} a''(-\xi) Z_3)(a(-\pi) - iK\beta^{-1} a'''(-\xi) a(\xi - \pi)) \\ &\quad - (Z_2 + K\beta^{-2} a'''(-\xi) Z_3)(\beta^{-2} a'''(-\pi)) \\ &\quad - iK\beta^{-1} a''(-\xi) a(\xi - \pi) \\ &= K\beta^{-2} (a''(-\xi) a(-\pi) - \beta^{-2} a'''(-\xi) a'''(-\pi)) Z_3 \\ &\quad + Z_1 (a(-\pi) - iK\beta^{-1} a'''(-\xi) a(\xi - \pi)) \\ &\quad - Z_2 (\beta^{-2} a'''(-\pi) - iK\beta^{-1} a''(-\xi) a(\xi - \pi)) \\ &= -K\beta^{-2} G(\pi) f'(\xi) + iK\beta^{-3} G(\pi) \int_{\pi}^{\xi} a''(\xi - y) F(y) dy \\ &\quad - \beta^{-2} \int_{\pi}^0 \{ a'(-y) (a(-\pi) - iK\beta^{-1} a'''(-\xi) a(\xi - \pi)) \\ &\quad - a''(-y) (\beta^{-2} a'''(-\pi) - iK\beta^{-1} a''(-\xi) a(\xi - \pi)) \} F(y) dy \\ &= -K\beta^{-2} G(\pi) f'(\xi) \\ &\quad - \beta^{-2} \int_{\pi}^0 (a'(-y) a(-\pi) - a''(-y) \beta^{-2} a'''(-\pi)) F(y) dy \\ &\quad + iK\beta^{-3} \int_{\xi}^0 (a'(-y) a'''(-\xi) - a''(-y) \\ &\quad \times a''(-\xi)) a(\xi - \pi) F(y) dy \\ &\quad + iK\beta^{-3} \int_{\pi}^{\xi} \{ G(\pi) a''(\xi - y) + (a'(-y) a'''(-\xi) - a''(-y) \\ &\quad \times a''(-\xi)) a(\xi - \pi) \} F(y) dy \\ &=: I_1 + I_2 + I_3 + I_4. \end{aligned}$$

By virtue of Lemma 2,

$$\begin{aligned} |G(\pi)| &= \frac{b^2}{4} \left| \cosh b(\pi - \xi) + \frac{e^{b\xi}}{2} (\cos b\pi - \sin b\pi) \right. \\ &\quad \left. - \frac{e^{b\pi}}{2} (\cos b\xi + \sin b\xi) + O(1) \right| \\ &\leq Cb^2 e^{b\pi}, \end{aligned}$$

hence

$$|I_1| \leq Cb^{-2} e^{b\pi} \|f\|_V.$$

Let $K(y) := a'(-y)a(-\pi) - a''(-y)\beta^{-2}a'''(-\pi)$ for $y \in [0, \pi]$. Then

$$I_2 = \beta^{-2} \int_0^{\pi} K(y)g(y) dy + i\beta^{-1} \int_0^{\pi} K(y)f(y) dy =: I_2^1 + I_2^2.$$

Since

$$\begin{aligned} K(y) &= \frac{b}{4} \{ (\sin by - \sinh by)(\cos b\pi + \cosh b\pi) \\ &\quad - (-\cos by + \cosh by)(-\sin b\pi - \sinh b\pi) \} \\ &= \frac{b}{4} \left\{ \cosh by \sinh b\pi - \sinh by \cosh b\pi \right. \\ &\quad - \frac{e^{by} - e^{-by}}{2} \cos b\pi + \frac{e^{b\pi} + e^{-b\pi}}{2} \sin by \\ &\quad - \frac{e^{b\pi} - e^{-b\pi}}{2} \cos by + \frac{e^{by} + e^{-by}}{2} \sin b\pi \\ &\quad \left. + \sin by \cos b\pi - \cos by \sin b\pi \right\} \\ &= \frac{b}{4} \left\{ \sinh b(\pi - y) + \frac{e^{b\pi}}{2} (\sin by - \cos by) \right. \\ &\quad \left. + \frac{e^{by}}{2} (-\cos b\pi + \sin b\pi) + O(1) \right\} \end{aligned}$$

we see that

$$\begin{aligned} |I_2^1| &\leq \beta^{-2} \|K(y)\|_H \|g\|_H \\ &\leq Cb^{-3} (\|\sinh b(\pi - y)\|_H + e^{b\pi} \|\sin by - \cos by\|_H \\ &\quad + \|e^{by}\|_H + O(1)) \|g\|_H \\ &\leq Cb^{-3} e^{b\pi} \|g\|_H. \end{aligned}$$

On the other hand, integrating by parts in I_2^2 yields

$$I_2^2 = i\beta^{-1} \left(- \int_0^{\pi} \tilde{K}(y) f'(y) dy + [\tilde{K}(y) f(y)]_0^{\pi} \right)$$

with

$$\begin{aligned} \tilde{K}(y) &= \int_0^y K(z) dz \\ &= \frac{1}{4} \left\{ -\cosh b(\pi - y) - \frac{e^{b\pi}}{2} (\cos by + \sin by) \right. \\ &\quad \left. + \frac{e^{by}}{2} (-\cos b\pi + \sin b\pi) + O(1) \right\}. \end{aligned}$$

Thus

$$|I_2^2| \leq C\beta^{-1} e^{b\pi} (\|f'\|_H + |f(\pi)|) \leq C\beta^{-1} e^{b\pi} \|f\|_V.$$

We now turn to the next term I_3 . We notice that

$$\begin{aligned} I_3 &= iK\beta^{-3} a(\xi - \pi) \int_{\xi}^0 (-H'(y))(g(y) + i\beta f(y)) dy \\ &= iK\beta^{-3} a(\xi - \pi) \int_0^{\xi} H'(y)g(y) dy - K\beta^{-2} a(\xi - \pi) \\ &\quad \times \int_0^{\xi} H'(y)f(y) dy =: I_3^1 + I_3^2. \end{aligned}$$

Since

$$\begin{aligned} H'(y) &= \frac{b^4}{4} \left\{ \cosh b(y - \xi) + \frac{e^{b\xi}}{2} (\sin by - \cos by) \right. \\ &\quad \left. - \frac{e^{by}}{2} (\cos b\xi + \sin b\xi) + O(b^2) \right\} \end{aligned}$$

we have that

$$\begin{aligned} |I_3^1| &\leq Cb^{-2} e^{b(\pi-\xi)} (\|\cosh b(y - \xi)\|_{L^2(0, \xi)} \\ &\quad + e^{b\xi} \|\sin by - \cos by\|_{L^2(0, \xi)} \\ &\quad + \|e^{by}\|_{L^2(0, \xi)} + O(b^2)) \|g\|_H \\ &\leq Cb^{-2} e^{b\pi} \|g\|_H. \end{aligned}$$

Integrating twice by parts in I_3^2 yields

$$\begin{aligned} I_3^2 &= -K\beta^{-2} a(\xi - \pi) \\ &\quad \times \left(\int_0^{\xi} G(y)f''(y) dy + [H(y)f(y) - G(y)f'(y)]_0^{\xi} \right) \end{aligned}$$

(since $G' = H$). Using (56) and the fact that $f(0) = f'(0) = 0$, we obtain that

$$\begin{aligned} |I_3^2 + K\beta^{-2} a(\xi - \pi)H(\xi)f(\xi)| &\leq C\beta^{-2} e^{b(\pi-\xi)} \\ &\quad \times (b^2 e^{b\xi} \|f''\|_H + b^2 e^{b\xi} |f'(\xi)|) \\ &\leq C\beta^{-1} e^{b\pi} \|f\|_V. \end{aligned}$$

Using (57) to estimate $H(\xi)$, we conclude that

$$I_3^2 = \frac{K}{16} b^{-1} e^{b\pi} (\cos b\xi + \sin b\xi)f(\xi) + O(\beta^{-1} e^{b\pi} \|f\|_V).$$

It remains to bound

$$\begin{aligned} I_4 &= iK\beta^{-3} \int_{\pi}^{\xi} L(y)(g(y) + i\beta f(y)) dy \\ &= iK\beta^{-3} \int_{\pi}^{\xi} L(y)g(y) dy - K\beta^{-2} \int_{\pi}^{\xi} L(y)f(y) dy \\ &=: I_4^1 + I_4^2 \end{aligned}$$

where we have set

$$L(y) := G(\pi)a''(\xi - y) - H'(y)a(\xi - \pi).$$

We first estimate the function L .

$$\begin{aligned} L(y) &= \frac{b^4}{8} \left\{ (\cosh b(\pi - \xi) + \frac{e^{b\xi}}{2} (\cos b\pi - \sin b\pi) \right. \\ &\quad - \frac{e^{b\pi}}{2} (\cos b\xi + \sin b\xi) + O(1)) \\ &\quad \times (-\cos b(\xi - y) + \cosh b(\xi - y)) \\ &\quad - (\cosh b(y - \xi) + \frac{e^{b\xi}}{2} (\sin by - \cos by) \\ &\quad - \frac{e^{by}}{2} (\cos b\xi + \sin b\xi) + O(b^2)) \\ &\quad \left. \times (\cos b(\xi - \pi) + \cosh b(\xi - \pi)) \right\} \\ &= \frac{b^4}{8} \left\{ e^{b\pi} \left(\frac{1}{2} (\cos b\xi + \sin b\xi) \cos b(\xi - y) \right. \right. \\ &\quad \left. \left. + \frac{1}{4} (\cos by - \sin by) \right) \right. \\ &\quad \left. + e^{by} \left(\frac{1}{2} (\cos b\xi + \sin b\xi) \cos b(\xi - \pi) \right. \right. \\ &\quad \left. \left. + \frac{1}{4} (\cos b\pi - \sin b\pi) \right) \right. \\ &\quad \left. + e^{b(\pi+\xi-y)} \left(-\frac{1}{4} \right) (\cos b\xi + \sin b\xi) + O(e^{b(\pi-\delta)}) \right\} \end{aligned}$$

for every $y \in [\xi, \pi]$ and some $\delta > 0$. Thus

$$\begin{aligned} |I_4^1| &\leq Cb^{-2} \|g\|_H \left(e^{b\pi} (\|\cos b(\xi - y)\|_{L^2(\xi, \pi)} \right. \\ &\quad \left. + \|\cos by - \sin by\|_{L^2(\xi, \pi)}) \right. \\ &\quad \left. + \|e^{by}\|_{L^2(\xi, \pi)} + \|e^{b(\pi+\xi-y)}\|_{L^2(\xi, \pi)} + O(e^{b(\pi-\delta)}) \right) \\ &\leq Cb^{-2} e^{b\pi} \|g\|_H. \end{aligned}$$

Next we introduce the function $\tilde{L}(y) := G(\pi)a(\xi - y) - G(y)a(\xi - \pi)$ which fulfills $\tilde{L}' = L$. Easy calculations show that

$$\begin{aligned} \tilde{L}(y) = \frac{b^2}{8} & \left\{ -e^{b\pi} \left(\frac{1}{2} (\cos b\xi + \sin b\xi) \cos b(\xi - y) \right. \right. \\ & \left. \left. + \frac{1}{4} (\cos by - \sin by) \right) \right. \\ & \left. + e^{by} \left(\frac{1}{2} (\cos b\xi + \sin b\xi) \cos b(\xi - \pi) \right. \right. \\ & \left. \left. + \frac{1}{4} (\cos b\pi - \sin b\pi) \right) \right. \\ & \left. + e^{b(\pi+\xi-y)} \left(-\frac{1}{4} \right) (\cos b\xi + \sin b\xi) + O(e^{b(\pi-\delta)}) \right\} \end{aligned}$$

and that

$$\begin{aligned} \tilde{L}'(y) = \frac{b^3}{8} & \left\{ e^{b\pi} \left(\frac{1}{2} (\cos b\xi + \sin b\xi) \sin b(y - \xi) \right. \right. \\ & \left. \left. + \frac{1}{4} (\cos by + \sin by) \right) \right. \\ & \left. + e^{by} \left(\frac{1}{2} (\cos b\xi + \sin b\xi) \cos b(\xi - \pi) \right. \right. \\ & \left. \left. + \frac{1}{4} (\cos b\pi - \sin b\pi) \right) \right. \\ & \left. - e^{b(\pi+\xi-y)} \left(-\frac{1}{4} \right) (\cos b\xi + \sin b\xi) + O(e^{b(\pi-\delta)}) \right\}, \end{aligned}$$

hence for all $y \in [\xi, \pi]$

$$|\tilde{L}(y)| \leq Cb^2 e^{b\pi} \quad \text{and} \quad |\tilde{L}'(y)| \leq Cb^3 e^{b\pi}.$$

Integrating twice by parts in I_4^2 , we obtain

$$I_4^2 = K\beta^{-2} \left(\int_{\xi}^{\pi} \tilde{L}(y) f''(y) dy + [\tilde{L}'(y)f(y) - \tilde{L}(y)f'(y)]_{\xi}^{\pi} \right),$$

hence

$$\begin{aligned} \left| I_4^2 - K\beta^{-2} [\tilde{L}'(y)f(y)]_{\xi}^{\pi} \right| & \leq C\beta^{-2} (b^2 e^{b\pi} \|f''\|_H \\ & \quad + b^2 e^{b\pi} (|f'(\pi)| + |f'(\xi)|)) \\ & \leq C\beta^{-1} e^{b\pi} \|f\|_V. \end{aligned}$$

On the other hand

$$\begin{aligned} K\beta^{-2} [\tilde{L}'(y)f(y)]_{\xi}^{\pi} & = \frac{K}{16} b^{-1} e^{b\pi} \{ -(\cos b\xi + \sin b\xi) f(\xi) \\ & \quad + c(b)f(\pi) + O(e^{-b\delta}) \|f\|_V \}, \end{aligned}$$

where the function $c(b)$ is defined in (60) and $\delta > 0$ is small enough. It follows that

$$I_3^2 + I_4^2 = \frac{K}{16} b^{-1} e^{b\pi} c(b) f(\pi) + O(\beta^{-1} e^{b\pi} \|f\|_V).$$

We conclude that

$$N = \frac{K}{16} b^{-1} e^{b\pi} c(b) f(\pi) + O(\beta^{-1} e^{b\pi} (\|f\|_V + \|g\|_H)) \quad (62)$$

and that (writing again the subscript n)

$$\begin{aligned} |\beta_n w_n(\pi)| & = \left| \beta_n \frac{N}{D} \right| \\ & \leq \left| \frac{(K/16) b_n e^{b_n \pi} c(b_n) f_n(\pi)}{D_1 + iD_2} \right| + C(\|f_n\|_V + \|g_n\|_H) \\ & \leq C(\|f_n\|_V + \|g_n\|_H) \\ & \rightarrow 0 \quad \text{as } n \rightarrow \infty \end{aligned}$$

by (28) and (58). Using (50), we obtain the promised contradiction to (36). The proof of Theorem 3 is achieved. \square

4.2 Polynomial decay rate

The next result asserts that a polynomial decay rate still holds true for almost every value of ξ .

Theorem 4: For almost every $\xi \in (0, \pi)$ we have for every $l > 1$

$$\sup_{|\beta| \geq 1} |\beta|^{-l} \|(i\beta - \mathcal{A})^{-1}\| < \infty. \quad (63)$$

This implies by Liu and Rao (2005) that for every positive integer k there exists a constant $C_k > 0$ such that

$$\|e^{t\mathcal{A}} z_0\|_{\mathcal{H}} \leq C_k \left(\frac{\ln t}{t} \right)^{k/l} (\ln t) \|z_0\|_{\mathcal{D}(\mathcal{A}^k)} \quad \forall z_0 \in \mathcal{D}(\mathcal{A}^k).$$

Proof of Theorem 4: Notice first that it is sufficient to prove (63) for all numbers $l \in (1, +\infty) \cap \mathbb{Q}$ and for a.e. $\xi \in (0, \pi)$. Pick any $l \in (1, +\infty) \cap \mathbb{Q}$, any $\xi \in (0, \pi)$ (to be specified later) and argue by contradiction as in the proof of Theorem 3. If (63) is false, then there exist $\beta_n \in (0, +\infty)$, $(w_n, v_n) \in \mathcal{D}(\mathcal{A})$ for $n = 1, 2, \dots$ such that

$$\|(w_n, v_n)\|_{\mathcal{H}} = 1, \quad \beta_n \rightarrow +\infty, \quad (64)$$

and such that, setting $(f_n, g_n) := (i\beta_n - \mathcal{A})(w_n, v_n)$,

$$\beta_n^l (f_n, g_n) \rightarrow (0, 0) \text{ in } \mathcal{H}. \quad (65)$$

Then (36) and (50) hold true, and to obtain a contradiction to (36) it is sufficient to prove that $\beta_n w_n(\pi) \rightarrow 0$. The same calculations as in the proof of Theorem 3 lead to $w(\pi) = N/D$ (the subscript n being again omitted), where N fulfills (62), and $D = D_1 + iD_2$ with D_1 and D_2 given by (55) and (58), respectively. We aim to prove that $|D| \geq C e^{b\pi}/b^{2+2\varepsilon}$ for any $\varepsilon > 0$, a.e. $\xi \in (0, \pi)$, and every $b > 0$ large enough. To this end, we notice first that

$$\begin{aligned} & (\cos b\xi + \sin b\xi)(\cos b(\xi - \pi) - \sin b(\xi - \pi)) \\ &= 2 \sin \pi \left(b \frac{\xi}{\pi} + \frac{1}{4} \right) \sin \pi \left(b \left(1 - \frac{\xi}{\pi} \right) + \frac{1}{4} \right). \end{aligned}$$

Let $\| |x| \| = \inf_{k \in \mathbb{Z}} |x - k|$. We need the following result, which is a variant of a classical result in Diophantine approximations (compare [Lang (1995, Theorem 4)]. Its proof may be found in Crépeau and Prieur (to appear).

Lemma 5: *Pick any $\varepsilon > 0$. Then for almost every $\alpha \in (0, 1)$, there is only a finite number of solutions $q \in \mathbb{N}^*$ to the inequality*

$$\left\| \left(q + \frac{1}{2} \right) \alpha + \frac{1}{4} \right\| < \frac{1}{q^{1+\varepsilon}}.$$

In particular, for a.e. $\alpha \in (0, 1)$ one may find a positive constant C such that for all $q \in \mathbb{N}^*$

$$\left\| \left(q + \frac{1}{2} \right) \alpha + \frac{1}{4} \right\| \geq \frac{C}{q^{1+\varepsilon}}.$$

Applying that property to ξ/π and to $1 - \xi/\pi$, we conclude that for a.e. $\xi \in (0, \pi)$, we have for some constant $C > 0$ and for all numbers $b = q + \frac{1}{2}$, $q \in \mathbb{N}^*$,

$$\left| \sin \pi \left(b \frac{\xi}{\pi} + \frac{1}{4} \right) \right| \geq \frac{C}{b^{1+\varepsilon}} \quad \text{and} \quad \left| \sin \pi \left(b \left(1 - \frac{\xi}{\pi} \right) + \frac{1}{4} \right) \right| \geq \frac{C}{b^{1+\varepsilon}},$$

hence $|c(b)| \geq C b^{-2-2\varepsilon}$ and

$$|D_2(b)| \geq C \frac{e^{b\pi}}{b^{1+2\varepsilon}}$$

if b is large enough. Since the function $c(b)$ is uniformly Lipschitz continuous on \mathbb{R} , the same inequalities hold (with different constants) for b (large enough) in

$$\bigcup_{q \in \mathbb{N}^*} \left(q + \frac{1}{2} - \frac{C'}{q^{2+2\varepsilon}}, q + \frac{1}{2} + \frac{C'}{q^{2+2\varepsilon}} \right),$$

if the constant $C' > 0$ is small enough. On the other hand, one may associate with that constant $C' > 0$ a constant $C'' > 0$ such that

$$|D_1(b)| \geq C'' \frac{e^{b\pi}}{b^{2+2\varepsilon}}$$

for b large enough fulfilling $|b - (q + \frac{1}{2})| \geq C'/q^{2+2\varepsilon}$ for each $q \in \mathbb{N}^*$. It follows that for a.e. $\xi \in (0, \pi)$, we have

$$|D(b)| \geq C \frac{e^{b\pi}}{b^{2+2\varepsilon}} \tag{66}$$

for b large enough. Gathering together (62) and (66), we arrive to

$$|\beta_n w_n(\pi)| \leq C b_n^{2+2\varepsilon} (\|g_n\|_H + \|f_n\|_V) \rightarrow 0$$

as $n \rightarrow +\infty$ by (65), if we pick $\varepsilon := l - 1$. Thus we have obtained the desired contradiction to (36). The proof of Theorem 4 is complete. \square

5. Conclusion

The paper was devoted to the output stabilization of a clamped-free beam with collocated piezoelectric sensor/actuator. Under the assumption that the actuator is touching the clamped extremity of the beam, it has been proved that the strong stability holds provided that the length ξ of the actuator does not belong to a certain dense countable set of $(0, \pi)$. Under this assumption it has been shown that the energy decreases exponentially if the ratio ξ/π belongs to a certain set of rational numbers. The question whether this last assumption may be dropped will be the purpose of further research in a near future.

Acknowledgements

The authors would like to thank Prof. Hisham Abou-Kandil, who brought the problem to their attention, and an anonymous referee for very useful comments. The second author was partially supported by HYCON Network of Excellence, contract number FP6-IST-511368, and the third author was partially supported by INRIA-CONICYT.

References

- K. Ammari and M. Tucsnak, "Stabilization of Bernoulli-Euler beams by means of a pointwise feedback force", *SIAM J. Control Optim.*, 39, pp. 1160–1181 (electronic), 2000.
- W. Arendt and C.J.K. Batty, "Tauberian theorems and stability of one-parameter semigroups", *Trans. Amer. Math. Soc.*, 306, pp. 837–852, 1988.
- H.T. Banks, W. Fang, R.J. Silcox and R.C. Smith, "Approximation methods for control of the acoustic/structure interaction with piezoceramic actuator", *J. Intelligent Material Syst. and Struct.*, 4, pp. 98–116, 1993.
- H.T. Banks and R.C. Smith, "The modelling of piezoceramic patch interactions with shells, plates and beams", *Tech. Report no. 92-66, Institute for Computer Applications in Science and Engineering*, Hampton, VA, 1992.
- G. Chen, S.G. Krantz, D.W. Ma, C.E. Wayne and H.H. West, "The Euler-Bernoulli beam equation with boundary energy dissipation", in *Operator methods for optimal control problems (New Orleans, La., 1986)*, volume 108 of *Lecture Notes in Pure and Appl. Math.*, pp. 67–96, New York: Dekker, 1987.
- F. Conrad, "Stabilization of beams by pointwise feedback control", *SIAM J. Cont. Optim.*, 28, pp. 423–437, 1990.
- E.F. Crawley and E.H. Anderson, "Detailed models for piezoceramics actuation of beams", *J. Intel. Mater. Syst. Struct.*, 1, pp. 79–83, 1990.
- E. Crépeau and C. Prieur, "Control of a clamped-free beam by a piezoelectric actuator", *ESAIM Control Optim. Calc. Var.*, 12, pp. 545–563, 2006.
- M.S. de Queiroz, D.M. Dawson, S.P. Nagarkatti and F. Zhang, *Lyapunov-based Control of Mechanical Systems*, Control Engineering, Boston, MA: Birkhäuser Boston Inc., 2000.
- P. Destuynder, "A mathematical analysis of a smart-beam which is equipped with piezoelectric actuators", *Control Cybernet.*, 28, pp. 503–530, 1999.
- P. Destuynder, I. Legrain, L. Castel and N. Richard, "Theoretical, numerical and experimental discussion on the use of piezoelectric devices for control-structure interaction", *European J. Mech. A Solids*, 11, pp. 181–213, 1992.
- F. Lun Huang, "Characteristic conditions for exponential stability of linear dynamical systems in Hilbert spaces", *Ann. Differential Equations*, 1, pp. 43–56, 1985.
- J. Lagnese and G. Leugering, "Uniform energy decay of a class of cantilevered nonlinear beams with nonlinear dissipation at the free end", in *Differential Equations with Applications in Biology, Physics, and Engineering (Leibnitz, 1989)*, volume 133 of *Lecture Notes in Pure and Appl. Math.*, pp. 227–239. New York: Dekker, 1991a.
- J.E. Lagnese and G. Leugering, "Uniform stabilization of a nonlinear beam by nonlinear boundary feedback", *J. Diff. Equa.*, 91, pp. 355–388, 1991b.
- S. Lang, *Introduction to Diophantine Approximations*, New York: Springer-Verlag, second edition, 1995.
- P. Le Gall, C. Prieur and L. Rosier, "Stabilization of a clamped-free beam with collocated piezoelectric sensor/actuator", *International Journal of Towography & Statistics*, to appear.
- J.-L. Lions and E. Magenes, *Non-homogeneous Boundary Value Problems and Applications*, Vol. I. Springer-Verlag, New York, Translated from the French by P. Kenneth, Die Grundlehren der mathematischen Wissenschaften, Band 181, 1972.
- Z. Liu and B. Rao, "Characterization of polynomial decay rate for the solution of linear evolution equation", *Z. Angew. Math. Phys.*, 56, pp. 630–644, 2005.
- Z. Liu and S. Zheng, *Semigroups associated with dissipative systems, volume 398 of Chapman & Hall/CRC Research Notes in Mathematics*. Boca Raton, FL: Chapman & Hall/CRC, 1999.
- Ö. Morgül, "Dynamic boundary control of an Euler-Bernoulli beam", *IEEE Trans. Automat. Cont.*, 37, pp. 639–642, 1992.
- J. Prüss, "On the spectrum of C_0 -semigroups", *Trans. Amer. Math. Soc.*, 284, pp. 847–857, 1984.
- R. Rebarber, "Exponential stability of coupled beams with dissipative joints: a frequency domain approach", *SIAM J. Cont. Optim.*, 33, pp. 1–28, 1995.
- M. Tucsnak, "Regularity and exact controllability for a beam with piezoelectric actuator", *SIAM J. Cont. Optim.*, 34, 922–930, 1996.